

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/SE05/000528

International filing date: 13 April 2005 (13.04.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/562,985  
Filing date: 19 April 2004 (19.04.2004)

Date of receipt at the International Bureau: 24 June 2005 (24.06.2005)


Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

21-04-2005

PA 1306843



# THE UNITED STATES OF AMERICA

**TO ALL TO WHOM THESE PRESENTS SHALL COME:**

**UNITED STATES DEPARTMENT OF COMMERCE**

**United States Patent and Trademark Office**

**April 12, 2005**

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.**

**APPLICATION NUMBER: 60/562,985**

**FILING DATE: April 19, 2004**

**By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS**



*M. Tarver*

**M. TARVER  
Certifying Officer**

13281 U.S. PTO

MAIL STOP PROVISIONAL APPLICATION  
PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c)

INVENTOR(S)

Given Name (first and middle [if any])	Family Name or Surname	Residence (City and State or Foreign Country)
Per	RYDBERG	Hagersten, Sweden

☐ Additional inventors are being named on the \_\_\_\_\_ separately numbered sheets attached hereto

TITLE OF THE INVENTION (500 characters max)

INVENTION OF A FLUORESCENT N-R-EDMAN PROCEDURE FOR ANALYSIS OF N-TERMINAL PROTEIN ADDUCTS WITH SPECTROPHOTMETRIC AND/OR MASS SPECTROMETRIC DETECTION

CORRESPONDENCE ADDRESS

Direct all Correspondence to:

☒ Customer Number

00466

Type Customer No. Here

Firm Name & Address:

Young & Thompson  
745 South 23<sup>rd</sup> Street  
Arlington, VA 22202  
USA

Telephone: 703-521-2297 - Facsimile: 703-685-0573

ENCLOSED APPLICATION PARTS (check all that apply)

<input checked="" type="checkbox"/> Specification	Number of Pages:	10	<input type="checkbox"/> CD(s), Number	
<input checked="" type="checkbox"/> Drawings	Number of Sheets:	22	<input checked="" type="checkbox"/> Other (specify)	
<input checked="" type="checkbox"/> Application Data Sheet			<u>Abstract</u>	

METHOD OF PAYMENT OF FILING FEES FOR PROVISIONAL APPLICATION FOR PATENT

- ☒ Applicant(s) claims small entity status.
- ☒ A check is enclosed in the amount of \$80.00 for the filing fee.
- ☒ The Commissioner is authorized to charge any underpayment or credit any overpayment to Deposit Account No. 25-0120

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

- ☒ No
- ☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

Docket No.: 1523-1010

By: Eric Jensen  
Eric Jensen, Reg. No. 37,855  
Telephone (703) 521-2297

Date: April 19, 2004

EJ/yr

PROVISIONAL APPLICATION FILING ONLY

Y&T April 19, 2004

# **Invention of a Fluorescent N-R-Edman procedure for analysis of N-terminal protein adducts with spectrophotometric and/or mass spectrometric detection**

## **Abstract**

We hereby introduce a novel method for analysis of electrophilic compounds measured *in vivo* as their corresponding adducts by means of the so-called "Fluorescent N-R-Edman procedure". This invention is based on the original observation that N-alkylated N-terminal protein adducts are detached with high selectivity from adducted proteins as their corresponding N-alkyl-valine phenylthiohydantoinines after derivatisation with phenyl isothiocyanate (PITC) or pentafluorophenyl isothiocyanate (PFPITC) in the so called N-alkyl Edman procedure. In the Fluorescent N-R-Edman procedure fluorescent Edman reagents are used (i.e. the isothiocyanates; FITC, DNITC and DABITC), detached as their corresponding thiohydantoinines which has been isolated from tested N-substituted amino acids model peptides and adducted protein. The analytes were separated on both liquid chromatography (LC) and capillary electrophoresis (CE) and detected by e.g. mass spectrometry (MS) and fluorescence spectroscopic techniques. Low limit of detection (LOD) was obtained using LC-MS/MS techniques (low fmol level on a standard instrument) and a potency for measurements down to low zmol ( $10^{-21}$ ) levels utilizing CE-laser induced fluorescence (LIF) analysis.

Due to the mild and non-discriminating conditions utilized for the Fluorescent N-R-Edman procedure its applicability range is wider than for all earlier existing methods for measurements of N-terminal protein adducts. This new method have excellent sensitivity and specificity, the range for measurement spans from adducts with low molecular masses to adducts that are thermo labile, have high molecular weight and/or have high polarity.

This new method should have the potency for routine analysis for hygienic surveillance, medical purpose and for analysis of forensic substances as their *in vivo* formed electrophilically reactive metabolites.

## **Background to the invention**

It has been demonstrated earlier that *in vivo* electrophilic compounds can be monitored by measuring the products (adducts) of their reaction with proteins, in particular hemoglobin

(Hb) (1-5). Important nucleophilic sites in Hb which are reactive under physiological conditions are the imidazole nitrogen atoms in histidine residues, sulfur atoms in cysteine and methionine residues, oxygen atoms in carboxyl groups and in hydroxyl groups in tyrosine and serine residues, and the  $\alpha$ -nitrogen atoms in the N-terminal valine residue of all four chains of human Hb (6).

The so called N-alkyl Edman procedure was developed for measurements of adducts (mainly low molecular weight adducts) to N-terminal valine residues in Hb (7). This method was based on the original Edman degradation procedure (8,9) used for protein sequencing. It was observed that N-terminal valine N-alkylated with a radioactively labelled 2-hydroxyethyl moiety from ethylene oxide was released spontaneously as a phenylthiohydantoin (PTH) under the conditions (pH >7) employed for the coupling reaction between phenyl isothiocyanate (PITC) and protein. The released PTH could be separated from unmodified N-terminal valine residues, as well as from the rest of the protein by extraction.

This observation led to the development of the N-alkyl Edman procedure for mass spectrometric (MS) quantitation of Hb adducts (10). Because of its usefulness, the N-alkyl Edman method has been applied in a number of laboratories for research purposes, dose monitoring and hygienic surveillance (11-16).

A brief description of the N-alkyl Edman procedure is presented in **Figure 1**. A sample of the globin (isolated from red blood cells by acid precipitation) is dissolved in formamide and pentafluorophenyl isothiocyanate (PFPITC) is then added, together with a small amount of aqueous 1 M NaOH in order to obtain a near neutral solution. The mixture is maintained at room temperature overnight, after which the temperature is raised to 45 °C for a couple of hours (17). The pentafluorophenylthiohydantoin (PFPTH) derivative of the terminal N-alkylvaline residues are released in high yield by this procedure and subsequently isolated by extraction.

Although the N-alkyl Edman procedure has become an established method for analysis of N-substituted haemoglobin adducts, the method is connected by limitations, e.g., the range of adducts that can be analysed. Small adducts, e.g., ethylene oxide and propylene oxide can be quantified at nmol/g globin level, which is sensitive enough for measurement of so called background adduct levels (levels without known exposure). However, adducts with a few polar groups are more difficult to measure, mainly because of elution problems in the gas chromatographic (GC) system prior to the MS detection. Some of these limitations can be solved, e.g., by further derivatisation (Paulsson et al), this approach is both time consuming and demands introduction of new steps to be developed for each specific adduct. Adducts

with; high molecular weight (>700 mass units), many polar groups and/or are thermo labile will be extremely difficult to analyse with the GC-MS based N-alkyl Edman procedure.

In order to overcome these problems and to optimize the sensitivity and increase the applicability range the "Fluorescent N-R-Edman procedure" has been developed. This invention is based on the principles of the original N-alkyl Edman procedure, alkylated N-terminals can be detached and measured as their corresponding thiohydantoin derivatives after coupling with isothiocyanate Edman reagents.

In contrast to the N-alkyl Edman procedure, this method is based on liquid chromatographic systems, e.g LC and capillary electrophoresis (CE) techniques with MS and/or fluorescence spectroscopic detection methods. Due to the mild and non-discriminating conditions utilized for the Fluorescent N-R-Edman procedure its applicability range is much wider compared to the N-alkyl Edman method. The introduction of fluorescent isothiocyanate reagents provides new possibilities for adduct measurements e.g. capillary electrophoresis with laser induced fluorescence detection (CE-LIF, ref) and HPLC with fluorescence detection for N-terminal protein adducts. The use of fluorescent isothiocyanates has been used for peptide sequencing (18,19) at nmolar levels by use of e.g., fluorescein isothiocyanate in combination by CE-LIF (21), in the study performed by Ireland *et al* the LOD were established to be on the low zmol level.

The Fluorescent N-R-Edman procedure is further improved by the utilization of ionisable groups in the applied reagents, e.g., tertiary amines and carboxyl groups, which provides possibilities to enrich the analytes on ion-exchangers, facilitates CE separation and also delivers higher affinity for ionisation in the MS, providing higher sensitivity. The principles of the Fluorescent N-R-Edman procedure is presented in Figure 2.

In order to compare the Fluorescent N-R-Edman procedure with the reagents that are used in the original N-alkyl Edman method and also recently has been applied for LC-MS/MS analysis (Ph.D., Hubert Vesber, using PFPITC, presented in Anaheim, CA May 2004, at the ACS meeting, acrylamide section), a comparative study were performed. The reagents for the N-alkyl Edman procedure, phenyl isothiocyanate (PITC) and pentafluorophenyl isothiocyanate (PFPITC) were evaluated towards three fluorescent isothiocyanate reagents; fluorescein isothiocyanate (FITC), 4-N,N-dimethylaminoazobenzene 4'-isothiocyanate (DABITC) and 4-dimethylamino-1-naphthyl isothiocyanate (DNITC) (see Figure 3). The tested reagents were reacted with valine and N-methyl valine, models for non-substituted N-terminal globin and N-substituted N-terminal valine in globine, N-(2-carbamoyl-ethyl)-valine were reacted with PFPITC. The formed analytes; phenyl-(valin and N-

methylvaline)thiohydantoin (PTH-Val and PTH-MeVal), pentafluorophenyl-*N*-(2-carbamoyl-ethyl)-valine thiohydantoin (PFPTH-AAVal), fluorescein-(valin and *N*-methylvaline)thiohydantoin (FTH-Val and FTH-MeVal), 4'-*N,N*-dimethylaminoazobenzene-4-(valin and *N*-methylvaline)thiohydantoin (DABTH-Val and DABTH-MeVal) and 4-dimethylamino-1-(valin and *N*-methylvaline)thiohydantoin (DANTH-Val and DANTH-MeVal) were then compared by measurements at various pH on UV, HPLC and LC-MS/MS. The fluorescent analytes were also measured on fluorescence spectroscopy (excitation and emission spectra) and on CE with diode array detection. As it was observed that one of the selected fluorescent reagent were superior in comparison with the rest of the reagents, this reagent were also evaluated on alkylated model peptides and globine adducted with acrylamide (AA), glycidamide (GA), propylene oxid (PE), cholesterol-5 $\alpha$ ,6 $\alpha$ -epoxide (Chol-EO) and 2-octadecyl-oxirane.

## Experimental

### Material and Methods

**Chemicals.** Structures of compounds 1-14 are given in Figure 4. Fluoresceine isothiocyanate (isomer I, <90%), pentafluorophenyl isothiocyanate (PFPITC) and phenyl isothiocyanate (PITC, purum) were obtained from Fluka. 4-*N,N*-Dimethylaminoazobenzene 4'-isothiocyanate (DABITC) and 4-dimethylamino-1-naphthyl isothiocyanate (DNITC) were obtained from Acros. Cholesterol-5 $\alpha$ ,6 $\alpha$ -epoxide, octadecyl-epoxide, L-Valine (Val), L-valinamide (ValNH<sub>2</sub>), *N*-methyl-*D,L*-valine (MeVal) and 5-isopropyl-3-phenyl-2-thiohydantoin (Val-PTH, 1) were obtained from Sigma. *N*-Methylvalylleucylanilide (MeValLeu-NH $\phi$  > 99 %) and valylleucylserine [ValLeuSer (H-Val-Leu-Ser-OH) 95%] were obtained from Bachem (Bubendorf, Switzerland). (<sup>2</sup>H<sub>3</sub>)Acetonitrile (99.8 % <sup>2</sup>H), (<sup>2</sup>H)chloroform (99.8 % <sup>2</sup>H), deuterium oxide (99.9 % <sup>2</sup>H, <sup>2</sup>H<sub>2</sub>O), and (<sup>2</sup>H<sub>4</sub>)methanol (99.8 % <sup>2</sup>H) were obtained from CIL (Andover, MA). 5-Isopropyl-1-methyl-3-phenyl-2-thiohydantoin (MeVal-PTH, 6), 5-isopropyl-3-pentafluorophenyl-2-thiohydantoin (Val-PFPTH, 12) and 5-isopropyl-1-methyl-3-pentafluorophenyl-2-thiohydantoin (MeVal-PFPTH, 13) were synthesised as described earlier (20). Glycidamide (GA) were synthesised. All other chemicals and solvents were of analytical grade.

**Instrumentation, methods for analysis and characterization.** <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a JEOL GSX 270 instrument at 270 MHz. All solvents used were fully deuterated, TMS was added as internal standard in chloroform, acetonitrile and methanol.

Methods and instrumentation for LC-MS/MS analysis. The LC-MS system comprised a Rheos 4000LC pump (Flux Instruments, Basel, Switzerland) interfaced with a LCQ (ThermoQuest, CA, USA). The MS was operated in electrospray ionisation (ESI) mode. The mobile phase consisted of 1:1, H<sub>2</sub>O:acetonitrile at an isocratic flow at 200 µL/min. The ion-source temperature was 120 °C, capillary voltage 3.5 kV and the cone voltage varied between 25 and 140 V. Nitrogen was used as drying gas at a flowrate of 250 L/h. Both the positive- and the negative-ion mode were used. MS-MS was performed by utilizing collision-induced dissociation (CID) of the [M+1] ion.

Studies of retention times of compounds 4-10 were carried out on HPLC, on a Shimadzu LC-4A connected with a Kromasil LC-18 column (250 x 10 mm) and a Shimadzu SPD-2AS detector ( $\lambda$  = 268 nm, D<sub>2</sub> lamp). Flow rate = 2.5 mL/min, loop 0.7 ml eluted with 2 % aqueous acetonitrile buffered with 0.02 % TFA.

TLC was performed using silica gel 60 f-254 plates (SiO<sub>2</sub>, Merck), spots were developed with UV and at long wave (378 nm) Melting points were determined on a Büchi 535 instrument. Mikrokemi AB, Uppsala, Sweden, performed elementary analyses. Measurements of pH were carried out on an Orion EA 920 pH-meter equipped with a Ross 8130 glass electrode.

The CE separation was performed on an HP 3D CE with a five channel diode array UV detector (Agilent, CA, USA). A fused silica capillary (i.d. 50 µm, o.d. 375 µm) with a total length of 64 cm and an effective length of 56 cm was used. The separation voltage was +30 kV, resulting in an separation current of 32 µA. The buffer system consisted of a 17 mM phosphate buffer (adjusted to pH 7) containing 20 mM SDS.

*Synthesis of N-Me(4-Me<sub>2</sub>N-Nafty)valin-tiohydantoin (5, DMAP-MeVal ).*

D,L-MeVal (prepared according to Rydberg *et al* 93, 93.0 mg, 0,708 mmol) was alkalinized with KOH (0,4 mmol) and dissolved in 0,5 M KHCO<sub>3</sub> (3 ml) and dioxan (2 ml). The solution was heated to 45 °C with magnetic stirring and p-dimethylnafthylisothiocyanate (DNITC, 91.3 mg, 0.40 mmol) dissolved in dioxane (2 ml) was added and the solution. The reaction was followed on TLC (SiO<sub>2</sub>, toluene and 2:1 toluene / ethyl acetate) with a blank reference (DNITC dissolved in the solvent mixture). After 60 min the reaction was completed, no DMNaf-ITC could be detected on TLC and the formed product gave fluorescence at long wave under the UV-lamp. The reaction was extracted with toluene and purified by column chromatography ((SiO<sub>2</sub>, toluene and 2:1 toluene / ethyl acetate), to yield 130 mg, 96 % yield. The product was recrystallized from ethanol/water (1:1), to yield white crystals (100 mg, 73 % yield), for NMR, see Figure 5.



*Synthesis of 4-Me<sub>2</sub>N-Naftyl)valin-thiohydantoin (4, DMAP-Val).* L-Valine (Sigma, 804 mg, 6.86 mmol) was alkalized with KOH (3.43 mmol) and dissolved in 0,5 M KHCO<sub>3</sub> (3 ml) and dioxan (3 ml). The solution was heated to 45 °C with magnetic stirring and DMNaf-ITC (88 mg, 0.385 mmol) dissolved in dioxane (2 ml) was added to the suspended solution. The reaction was followed on TLC (SiO<sub>2</sub>, Ethyl acetate / MeOH 4:1, toluene and 2:1 toluene / ethyl acetate) with a reference without addition of Valine. After 60 min the reaction was in principal completed, no DMNaf-ITC could be detected on TLC and the formed product, the 4-Me<sub>2</sub>N-Naftyl)-tiocarbamoyl-valinate gave fluorecence at long wave under the UV-lamp (tailing spot on TLC eluted with Ethyl acetate / MeOH 4:1). The reaction was acidified with hydrochloric acid (3 ml conc.) and the cyclisation was interrupted after 2 h, at 45 ° by neutralization with solid KHCO<sub>3</sub> until no more carbodioxide was formed. The reaction mixture was diluted with water (100 ml) and the product was purified by extraction with CHCl<sub>3</sub> (125 ml). The organic phase was dried with Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated to yield 126 mg, yield 0.96 % as a white solid.

The syntheses of compound 6-14 were performed as described above, the products were confirmed by <sup>1</sup>H and <sup>13</sup>C NMR spectra and by LC-MS/MS (ESI) positive and negative ionization, eg., **Figure 6**, <sup>13</sup>C NMR spectra of FTH-MeVal, **Figure 7**, <sup>1</sup>H NMR spectra of FTH-MeVal and **Figure 8**, <sup>13</sup>C NMR spectra of FTH -MeVal.

### Results and discussion

In order to evaluate the potency of the tested isothiocyanates the relative response of compound 1-14 (see **Figure 3**) were measured and compared using fluorescence spectroscopy (compound 4-10), HPLC, CE and UV spectroscopy. For the main part of these studies pH were alternated below and above the pKa for the respective analyte (for pKa values see **Figure 4**). The studies on relative fluorescence were performed on the fluorescent analytes (compound 4-10 in **Figure 4**) by measurements of excitation and emission spectra. The results from this study is presented in **figure 9-12**, FTH-Val and FTH-MeVal gave both in principle identically and high response at pH above 5, showing that the adduct, e.g., the methyl in FPT-MeVal (**9**) does not effect the spectroscopic properties in comparison with FTH-Val (**8**). In these studies FTH-Val and FTH-Me Val gave superior response compared to the other used reagents.

In order to evaluate the potency of the tested isothiocyanates the relative response of compound 1-14 (see Figure 3) were measured and compared using UV spectroscopy, HPLC, CE and fluorescence spectroscopy (compound 4-10). For the main part of these studies pH were alternated below and above the pKa for the respective analyte (for pKa values see Figure 4). The studies on relative fluorescence were performed on the fluorescent analytes by measurements of excitation and emission spectra. The results from this study is presented in figure 9-12, FTH-Val and FTH-MeVal gave both in principle identically and high response at pH above 5, showing that the adduct, e.g., the methyl in FPT-MeVal (9) does not effect the spectroscopic properties in comparison with FTH-Val (8). In these studies FTH-Val and FTH-Me Val gave superior response compared to the other used reagents.

In order to compare the relative sensitivity on LC-MS/MS compound 1-9 were compared by injecting 5  $\mu$ l directly in the electrospray interface of the TSQ (n=3 for each compound), the cone voltage were adjusted in order to obtain maximum sensitivity for the each specific analyte. The LC-MS/MS(ESI) measurements were performed using well adopted buffer systems, e.g., 0.1 % TFA, 0.1 % ammonium acetate and 0.3 mM aqueous ammonia. For LOD measurements the flow rate were 200  $\mu$ l/min of the aqueous buffer/acetonitrile [1:1 (v/v)] the analytes were dissolved in the same buffers and solvent mixtures at concentrations ranging from 10  $\mu$ g/ml down to 1 ng/ml depending on their response for each used condition. The results from this study are summarized in Figures 14-20 and in Table 1. As compound 8 and 9, FTH-Val and FTH-MeVal, gave superior response under the tested conditions for the LC-MS analysis in combination with excellent column separation on HPLC ( $C_{18}$ -columns, especially when eluted at acidic conditions), they were selected for further studies. In order to investigate the applicability range for this reagent, valine and globine were adducted with glycidamide, propylene oxide and the high molecular weight adducts cholesterol-5 $\alpha$ ,6 $\alpha$ -epoxide and octadecyl-epoxide. The formed thiohydantoinnes (compounds 10-14) were characterized on LC-MS/MS and the adducted globins could be measured after derivatisation (90 min in 0.5 M aqueous  $KHCO_3$ /2-propanol [2/1 (v/v)]), followed by size discriminating ultra filtration (MWCO 5000), evaporation filtrate or concentration of the analytes from the filtrate by use of anion exchangers. This approach is not only time saving but is also non-discriminating especially for analysis of adducts with high polarity as they often are lost or gives poor yields on liquid/liquid or solid phase extraction clean up steps.

In order to evaluate the possibility to measure N-terminal adducts with such sensitive techniques as CE-LIF, the separation between F-MeValTH and F-ValTH on CE were evaluated. A complete baseline separation was obtained using the stated conditions e.g., in 17 mM phosphare buffer (adjusted to pH 7) containing 20 mM SDS (see Figure 21). The intensity of the fluorescence reaches an optimum around this pH for both FTH-Val and FTH-MeVal, which opens up the possibility to utilize

CE-LIF for adducts measurements down to the low zmol ( $10^{-21}$ ) levels, which has been shown by Ireland *et al.* (21). In their study 18 of 20 coded FTH-amino acids were separated and analysed with LOD of around 10 zmol. Their results are truly encouraging for miniaturizing the fluorescent-N-R Edman procedure, aiming to measure adduct spectra from a few  $\mu$ l blood, easily available by a prick in the fingertip.

## References

1. Ehrenberg, L., Hiesche, K. D., Osterman-Golkar, S. and Wennberg, I. (1974) Evaluation of genetic risks of alkylating agents: Tissue doses in the mouse from air contaminated with ethylene oxide. *Mutat. Res.* **24**, 83-103.
2. Osterman-Golkar, S., Ehrenberg, L., Segerbäck, D. and Hällström, I. (1976) Evaluation of genetic risks of alkylating agents. II. Haemoglobin as a dose monitor. *Mutat. Res.* **34**, 1-10.
3. Neumann, H. G., Baur, H. and Wirsing, R. (1980) Dose-response relationship in the primary lesion of strong electrophilic carcinogens. *Arch. Toxicol. Suppl.* **3**, 69-77.
4. Skipper, P. L., and Tannenbaum, S. R. (1990) Protein adducts in the molecular dosimetry of chemical carcinogens. *Carcinogenesis* **11**, 507-518.
5. Törnqvist, M., Fred, C., Haglund, J., Helleberg, H., Paulsson, B. and Rydberg, P. (2002) Protein adducts: quantitative and qualitative aspects of their formation, analysis and applications. *J. Chromatogr. B* **778**, 279-308.
6. Osterman-Golkar, S. and Ehrenberg, L. (1980) Alkylation of macromolecules for detecting mutagenic agents. *Teratog. Carcinog. Mutag.* **1**, 105-127.
7. Jensen, S., Törnqvist, M., and Ehrenberg, L. (1984) Hemoglobin as a dose monitor of alkylating agents: Determination of alkylation products of N-terminal valine. (F.J.de Serres and R.W.Pero, Ed.) pp 315-320, Plenum Press, New York
8. Edman, P. (1950) Methods for determination of the amino acid sequence in peptides. *Acta Chem. Scand.* **4**, 283-298.
9. Edman, P., and Henschen, A. (1975) Sequence determination. In *Protein Sequence Determination* (Needleman S. B., Ed.) pp 232-279, Springer-Verlag, Berlin
10. Törnqvist, M., Mowrer, J., Jensen, S. and Ehrenberg, L. (1986) Monitoring of environmental cancer initiators through hemoglobin adducts by a modified Edman method. *Anal. Biochem.* **154**, 255-266.
11. Tate, A. D., Grummt, T., Törnqvist, M., Farmer, P. B., van Dam, F. J., van Mossel, H., Schoemaker, H. M., Osterman-Golkar, S., Uebel, C., Tang, Y. S., Zwinderman, A. H., Natarajan, A. T. and Ehrenberg, L. (1991) Biological and chemical monitoring of occupational exposure to ethylene oxide. *Mutat. Res.* **250**, 483-497.
12. Törnqvist, M., Magnusson, A. L., Farmer, P. B., Tang, Y. S., Jeffery, A. M., Wazneh, L., Beulink, G. D. T., van der Waal, H. and van Sittert, N. J. (1992) Ring test for low-levels of N-(2-hydroxyethyl)valine in human hemoglobin. *Anal. Biochem.* **203**, 357-360.
13. Bergmark, E., CALLEMAN, C. J., He, F., and Costa, L. G. (1993) Hemoglobin adducts in humans occupationally exposed to acrylamide. *Toxicol. Appl. Pharmacol.* **120**, 45-54.
14. Fidder, A., Noort, D., de Jong, A. L., Trap, H. C., de Jong, L. P., and Benschop, H. P. (1996) Monitoring of in vitro and in vivo exposure to sulfur mustard by GC/MS determination of the N-terminal valine adduct in Hb after a modified Edman degradation. *Chem. Res. Toxicol.* **9**, 788-792.
15. Thier, R., Lewalter, J., Kempkes, M., Selinski, S., Bruning, T., and Bolt, H. M. (1999) Haemoglobin adducts of acrylonitrile and ethylene oxide in acrylonitrile workers, dependent of polymorphisms of the glutathione transferases GSTT1 and GSTM1. *Occup. Environ. Med.* **56**, 197-202.
16. Begemann, P., Sram, R. J. and Neumann, H. G. (2001) Hemoglobin adducts of epoxybutene in workers occupationally exposed to 1,3-butadiene. *Arch. Toxicol.* **74**, 680-687.
17. Törnqvist, M. (1994) Epoxide adducts to N-terminal valine of hemoglobin. *Methods Enzymol.* (Everse J., Vandergriff K. D., and Winslow R. W., Eds.) pp 650-657, Academic Press, New York

18. Edman, P. (1950) Methods for determination of the amino acid sequence in peptides. *Acta Chem. Scand.* **4**, 283-298.
19. Edman, P., and Henschen, A. (1975) Chapter Title (Needleman, S. B.) pp Chap 8, Springer-Verlag, Berlin and New York
20. Rydberg, P., Luning, B., Wachtmeister, C. A. and Törnqvist, M. (1993) Synthesis and characterization of *N*-substituted valines and their phenyl-thiohydantoins and pentafluorophenyl- thiohydantoins. *Acta Chem. Scand.* **47**, 813-817.
21. Ireland, I. D., Lewis, D. F., Li, X., Renborg, A., Kwong, S., Chen, M., and Dovichi, N. J. (1997) Double Coupling Edman Chemistry for High-sensitivity Automated Protein Sequencing. *J. Protein Chem.* **16**, 491-493.


# 1. Tables

**Table 1.** Comparison of relative sensitivities obtained by measurements of limit of detection detection LOD on LC-MS/MS(ESI), using various pH modifiers/buffers. LOD were measured on positive and negative ions with direct injections (5 ul loop). The lowest obtained sensitivity is set to one, and the others are set relative to FTH-MeVal in 0.3 mM NH<sub>3</sub>, negative ions measured (marked yellow), this where calculated to be 2,6 fmol.


substance	Positive 0,1% NH <sub>4</sub> oAc	Positive 0,1% TFA	Positive 0,3mM NH <sub>3</sub>	Positive Without buffer	Negative 0,1% NH <sub>4</sub> oAc	negative 0,3mM NH <sub>3</sub>	negative Without buffer
FTH-MeVal	15,8	3,9	4,6	447,6	13,8	1,0	50,1
FTH-Val	3,6	1,1	n.d	393,1	1,6	n.d	57,3
DABTH-MeVal	60,4	26,9	9,8	377,7	45,1	55,9	386,7
PTH-MeVal	14439,5	7821,4	594,1	36098,9	6618,1	902,5	895,3
NTH-MeVal	940,8	90,9	460,7	932,1	216593,1	15752,2	984,8

Figures:


**Figure 1.** Principles of the N-alkyl Edman procedure.



**Figure 2. Principles of the Fluorescent N-R-Edman procedure.**




**Figure 3. Used isothiocyanate reagents in this study**







**Figure 4. Structures, abbreviations and molecular weights of the studied compounds**



Substance 1, R=H, PTH-Val, M=234.31 g/mol  
 Substance 2, R=Me, PTH-MeVal, M=248.35 g/mol  
 (non-fluorescent analytes)




Substance 3, R= CH<sub>2</sub>CH<sub>2</sub>CONH<sub>2</sub>, PTH-AAVal, M=395.35 g/mol  
 (non-fluorescent analyte)




Substance 4, R=H, DMAP-Val, M=327.44  
 Substance 5, R=Me, DMAP-MeVal, M=341.47  
 (fluorescent analytes, pK<sub>a</sub> for the dimethylamine amine calculated to be 6, (±0.3))

TITLE: INVENTION OF A FLUORESCENT N-R-EDMAN PROCEDURE FOR ANALYSIS OF N-TERMINAL PROTEIN ADDUCTS WITH SPECTROPHOTOMETRIC AND/OR MASS SPECTROMETRIC DETECTION



Substance 6, R=H, DABTH-Val M=381.50  
 Substance 7, R=Me DABTH-MeVal M=395.52  
 (fluorescent analytes, pKa for the dimethylamine amine calculated to be 3.4, ( $\pm 0.3$ ))



Substance 8, R=H FTH-Val M=488.51  
 Substance 9, R=Me FTH-MeVal M=502.54  
 Substance 10, R=CH<sub>2</sub>CH<sub>2</sub>CONH<sub>2</sub> FTH-AAVal M=559.59  
 Substance 11, R=CH<sub>2</sub>CH(OH)CONH<sub>2</sub> FTH-GAVal M=575.59  
 Substance 12, R=CH<sub>2</sub>CH[O]CH<sub>2</sub> FTH-POVal M=546.59  
 Substance 13, R=CH<sub>2</sub>CH[O]CH[CH<sub>2</sub>(n=15)]CH<sub>3</sub>, FTH-C<sub>18</sub>EO M=756.99  
 Substance 14, R=5a or 6a adduct to cholesterol epoxide: FTH-Chol-O M=891.17  
 (fluorescent analytes, pKa for the carboxyl group calculated to be 3.27 ( $\pm 0.3$ ))

Figure 5, <sup>13</sup>C NMR spectra of DMAP-MeVal

**TITLE: INVENTION OF A FLUORESCENT N-R-EDMAN PROCEDURE FOR ANALYSIS OF N-TERMINAL PROTEIN ADDUCTS WITH SPECTROPHOTOMETRIC AND/OR MASS SPECTROMETRIC DETECTION**





Figure 6,  $^{13}\text{C}$  NMR spectra of FTH-MeVal



**Figure 7,  $^1\text{H}$  NMR spectra of FTH-MeVal**







Figure 8,  $^{13}\text{C}$  NMR spectra of FTH -MeVal



**Figure 9.** Fluorescens measurements; the excitation and emission spectra of FTH-Val in dry acetonitrile at various concentrations.




**Figure 10.** Fluorescens measurements; the excitation and emission spectra of FTH-MeVal in at pH 1, 4, 7 and 9 at 0.5  $\mu\text{g/ml}$ .







**Figure 11.** Fluorescens measurements; the excitation and emission spectra of DABTH-MeVal in at pH 1, 4 7 and 9 at 1  $\mu$ g/ml.




**Figure 12.** Comparative studies of the excitation and emission spectra of the fluorescent analytes FTH-MeVal, FTH-Val, DABTH-MeVal, DABTH-Val, DNTH-MeVal, DNTH-Val and the reference fluorantene at the given concentrations in dry acetonitrile



**Figure 13.** Fluorescens measurements; the excitation and emission spectra of DAMTH-Val at pH 1, 4 7 and 9 at 1  $\mu\text{g/ml}$ .




**Figure 14.** Comparison between the tested analytes by measurements of limit of detection LOD on LC-MS/MS with electrospray ionisation and a 5 ul loop, without buffers, direct injections.




Substans	Precurs or ion (m/z)	Product ion (m/z)	Collision energi (rel. %)
PTH Me Val -	247.0	232.0	40
HOEtVal-PFPTH	367.1	347.1	25
FTH-MeVal -	501.2	457.3	35
FTH-Val -	487.2	443.3	30
DABTH-MeVal -	487.2	379.3	40
DABTH-Val -	380.2	264.3	35
NafTH-MeVal -	340.2	325.2	35
NafTH-Val -	326.1	210.2	35
PTH Me Val +	249.2	176.0	35
FTH-MeVal +	503.3	460.1	40
FTH-Val +	489.2	390.3	40
DABTH-MeVal +	396.2	219.0	40
DABTH-Val +	382.2	205.0	40
NafTH-MeVal +	342.2	327.1	35
NafTH-Val +	328.2	219.0	40


**Figure 15.** Comparison between the tested analytes by measurements of limit of detection LOD on LC-MS/MS with electrospray ionisation and a 5 ul loop, with an addition of 0.3 mM ammonia (pH around 9) with direct injection.




**Figure 16.** Comparison between the tested analytes by measurements of limit of detection LOD on LC-MS/MS with electrospray ionisation and a 5 ul loop, with an addition of 0.1 % TFA (pH around 1) with direct injection.



**Figure 17.** Comparison between the tested analytes by measurements of limit of detection LOD on LC-MS/MS with electrospray and a 5 ul loop, with additions of specified buffers, measured on positive ions with direct injections.




**Figure 18.** Comparison between the tested analytes by measurements of limit of detection LOD on LC-MS/MS with electrospray and a 5 ul loop, with additions of 0.2 % ammonium acetate (pH 6.96) , measured on negative ions with direct injections.






**Figure 19.** Comparison between the tested analytes by measurements of the limit of detection LOD on LC-MS/MS with electrospray and a 5 ul loop, with additions of 0.3 % ammonia (pH around 9) , measured on negative ions with direct injections.



**Figure 20.** Comparison between the tested analytes by measurements of the limit of detection LOD on LC-MS/MS with electrospray and a 5 ul lop, with additions of various pH modifying buffers , measured on negative ions with direct injections




DOCKET No.: 1523-1010  
INVENTOR: PER RYDBERG  
FILING DATE: APRIL 19, 2004

TITLE: INVENTION OF A FLUORESCENT N-R-EDMAN PROCEDURE FOR ANALYSIS OF N-TERMINAL PROTEIN ADDUCTS WITH  
SPECTROPHOTOMETRIC AND/OR MASS SPECTROMETRIC DETECTION

**Figure 21.** Separation between FTH-MeVal and FTH-Val on Capillary electrophoresis measured with diode array detection at 220 nm.

FTH-Val elutes at 7.44 min and FTH MeVal elutes at 8.91 min (phosphate buffer 17 mM, pH 7, 20 mM SDS). 30 kV, 52 cm capillary, 1 nl injected.



Application Data Sheet

**Application Information**

Application Type::	Provisional
Subject Matter::	Utility
Suggested Classification::	
Suggested Group Art Unit::	
CD-ROM or CD-R?::	None
Number of CD disks::	
Number of Copies of CDs::	
Sequence Submission?::	None
Computer Readable Form (CRF)::	No
Number of copies of CRF::	0
Title::	INVENTION OF A FLUORESCENT N-R- EDMAN PROCEDURE FOR ANALYSIS OF N-TERMINAL PROTEIN ADDUCTS WITH SPECTROPHOTMETRIC AND/OR MASS SPECTROMETRIC DETECTION
Attorney Docket Number::	1523-1010
Request for Early Publication?::	No
Request for Non-Publication?::	No
Suggested Drawing Figure::	
Total Drawing Sheets::	23
Small Entity?::	Yes
Latin Name::	
Variety Denomination Name::	
Petition Included?::	No
Petition Type::	
Licensed US Gov't Agency::	
Contract or Grant Numbers::	
Secrecy Order in Parent	No
Appl.?::	

**Applicant Information**

Applicant Authority Type:: Inventor  
Primary Citizenship Country:: SWEDEN  
Status:: Full Capacity  
Given Name:: PER  
Middle Name::  
Family Name:: RYDBERG  
Name Suffix::  
City of Residence:: HAGERSTEN  
State or Province of  
Residence::  
Country of Residence:: SWEDEN  
Street of Mailing TOMTRATTSVAGEN 37  
Address::  
City of Mailing Address:: HAGERSTEN  
State or Province of Mailing Address::  
Country of Mailing Address:: SWEDEN  
Postal or Zip Code of Mailing Address:: SE-129 31

**Correspondence Information**

Correspondence Customer 000466  
Number::

**Representative Information**

Representative Customer	000466
Number::	

**Domestic Priority Information**

Application::	Continuity Type::	Parent Application::	Parent Filing Date::

**Foreign Priority Information**

Country::	Application Number::	Filing Date::	Priority Claimed::

**Assignment Information**

Assignee Name::

Street of Mailing

Address::

City of Mailing Address::

State or Province of Mailing Address::

Country of Mailing Address::

Postal or Zip Code of Mailing Address::